

NECHAYEV, G.A.; MAKASHEV, S.D.

Extending the raw material base. TSement 28 no.2:3-5
Mr-Apr '62. (MIRA 15:8)

1. Ministerstvo geologii i okhrany nedr SSSR.
(Cement industries)

MAKASHEV, S. D.

ONUFRIYEV, I.A., red.; BARSKOV, I.M., red.; DMITRIYEV, A.D., red.;
MAKASHEV, S.D., red.; PEVZNER, A.S., red. izd-va; GILSONSON,
F.G., techn. red.

[Abridged stenographic report of the All-Union Conference on Construction. Moscow, 1958] Sokrashchennyi stenograficheskiy otchet. Moskva, Gos. izd-vo lit-ry po stroit., arkhitekt. i stroit. materialam, 1958. 334 p. (MIRA 11:10)

1. Vsesoyuznoye soveshchaniye po stroitel'stvu. Moscow, 1958. (Construction industry--Congresses)

MAKASHEV, M.Kh.; TKHORZHEVSKIY, O.A.

Aging of high-Q AT-cut quartz resonators. Izv.tekh. no.5:53-54
My '63. (MIRA 16:10)

MAKASHEV, R.K.; KAZACHENKO, I.V.

Distribution and excretion of radioactive calcium from the organism of healthy and lead-poisoned animals. Izv. AN Kazakh. SSR. Ser. med. nauk no.1850-55 '62 (MIRA 17:7)

MAKASHEV, K.K.; RAKHIMOVA, Z.P.

Distribution and discharge of P^{32} from the organism of healthy
and lead-poisoned animals. Izv. AN Kazakh. SSR. Ser. med. nauk
11 no.2:44-49 '64. (MIRA 17:7)

MAKASHTEV, K.K.;SHESTAKOVA, N.P.

Distribution and discharge of lead from the organism of healthy
animals and those poisoned by lead. Trudy Inst. kraev. pat AN
Kazakh. SSR 9:129-135'61. (MIRA 16:7)
(LEAD POISONING) (EXCRETION)

MAKASHEV, K.K.; AKHMEDOVA, A.S.

Effect of calcium and disodium salt of ethylenediaminetetraacetic acid and cortisone on the distribution of phosphorus and calcium in organs and tissues and their removal from the body in lead. Trudy Inst.kraev.pat. AN Kazakh.SSR 10:190-197 '62.

(MIRA 1685)

(LEAD POISONING) (ACETIC ACID—THERAPEUTIC USE)
(CORTISONE)

MAKASHEV, K.K.

Effect of calcium and disodium salt of ethylenediaminetetraacetic acid on the absorption and accumulation of lead by and its removal from the body in lead poisoning. Trudy Inst.kraev.pat. AN Kazakh. SSR 10:180-189 '62. (MIRA 16:5)
(LEAD POISONING) (ACETIC ACID—THERAPEUTIC USE)

BUTRIMOVA, N.P.; MAKASHEV, K.K.

Effect of sodium salicylate on the development of experimental
silicosis in white rats. Trudy Inst.kraev.pat. AN Kazakh. SSR
10:78-94 '62. (MIRA 16:5)
(LUNGS--DUST DISEASES) (SODIUM SALICYLATE--THERAPEUTIC USE)

ATCHABAYOV, B.A., kand.med.nauk; MAKASHEV, K.K., kand.med.nauk; SHESTAKOVA,
N.P.

Fate of lead introduced into the organism. Vest.AN Kazakh.SSR 17
no.5:48-55 My '61. (MIRA 14:6)
(LEAD IN THE BODY)

MAKASHEV, K.K.

Changes in the body's immunological properties in lead poisoning.
Trudy Inst.kraev.pat. AN Kazakh.SSR 4:34-41 '56. (MLRA 10:3)
(LEAD POISONING) (IMMUNITY)

MAKASHEV, K.K.

ATCHARAROV, B.A.; MAKASHEV, K.K.

Behavior of lead in the body. Trudy Inst.kraev.pat. AN Kazakh, SSR
4:5-21 '56. (MLRA 10:3)
(LEAD IN THE BODY)

MAKASHEV, K.K.

Changes in complement and immune antibodies titer during complete
replacement and cross transfusion of blood. Trudy Inst.kraev.pat.
AN Kazakh.SSR 2:32-59 '54. (MLRA 10:1)
(BLOOD--TRANSFUSION) (COMPLEMENTS (IMMUNITY))
(ANTIGENS AND ANTIBODIES)

MAKASHEV, I.K.

Changes in the biological properties of blood in experimental
morphine shock. Trudy Inst.kraev.pat. AN Kazakh.SSR 1:140-143 '52.
(BLOOD--ANALYSIS AND CHEMISTRY) (MIRA 10:2)
(MORPHINE) (SHOCK)

GLOZMAN, O.S.; MAKASHEV, K.K.

Structure method in determining the toxin content of solutions and
the degree of toxemia. Izv. AN Kazakh.SSR. Ser.kraev.pat. no.6:
147-151 '50. (MLRA 9:8)

(TOXINS AND ANTITOXINS)

MAKASHEV, K. A.

The second prize (imeni N. A. Minkevich) was awarded to Candidate of Technical Sciences V. A. Yakovlev, Engineers Ya. N. Spektor and K. A. Makashev for the paper "New Heat Treatment Technology for Tubular Components of a Complex Geometrical Shape Using Induction Heating for the Hardening Process".

Results of the 1958 Competition for Obtaining imeni D. K. Chernov and imeni N. A. Minkevich Prizes, Metallovedeniye i termicheskaya obrabotka metallov, 1959, No. 6, pp 62-64

MAKASHEV, A.P., prof.; POLETAYEVA, N.N., starshiy nauchnyy sotrudnik; ISA-
GULYAN, E.A., mladshiy nauchnyy sotrudnik

Experimental storage of apples in film wrapping material and con-
tainers. Khol.tekh. 41 no.1:36-41 Ja-F '64. (MIRA 17:3)

1. Krasnodarskiy nauchno-issledovatel'skiy institut pishchevoy pro-
myshlennosti.

MAKASHEV, A.P.; Prinimali uchastiye: ALDAKIMOVA, A.Ya.; MINKINA, A.I.,
mladshiy nauchnyy sotrudnik; SOKOLOVA, Ye.V.

[Use of carbon dioxide in fish preservation]. Primenenie ugle-
kisloty pri khraneni ryby. Moskva, Pishchepromizdat, 1959. 136 p.
(Moscow. Vsesoiuznyi nauchno-issledovatel'skii institut morskogo
rybnovo khoziaistva i okeanografii. Trudy, vol. 37). Trudy VNIRO
37 '59. (MIRA 17:4)

1. Starshiye laboranty tekhnologicheskoy laboratorii Dono-Kuban-
skogo otdeleniya Azovsko-Chernomorskogo nauchno-issledovatel'skogo
instituta morskogo rybnogo khozyaystva i okeanografii (for Aldaki-
mova, Sokolova). 2. Tekhnologicheskaya laboratoriya Dono-Kubansko-
go otdeleniya Azovsko-Chernomorskogo nauchno-issledovatel'skogo
instituta morskogo rybnogo khozyaystva i okeanografii (for Min-
kina).

5(1)

AUTHOR: Makashev, A. P.

SOV/32-25-3-55/62

TITLE: Gas Analyzer - Pipette (Gazoanalizator - pipetka)

PERIODICAL: Zavodskaya Laboratoriya, 1959, Vol 25, Nr 3, pp 378-379 (USSR)

ABSTRACT: The apparatus described was designed for the determination of carbon dioxide, nitrogen, and carbon monoxide gases. Accuracy of measurement at a small consumption of reagents and an analysis duration of 1-1.5 minutes is given as 0.2-0.5 %. The apparatus (Fig) has a graduated pipette with a length of about 400 mm and a diameter of about 1 mm. A small diaphragm pump is attached to the upper end of the pipette; when a screwhead is turned, the pump sucks the gas or liquid sample into the pipette. The absorption liquid is introduced in the same manner into the pipette filled with the gas sample, and gas and absorption liquid are mixed by turning the screwhead back and forth. Comparative analyses carried out in the apparatus described and in the apparatus according to Orsat-Fischer (CO₂ determinations)(Table) showed practically equal results. There are 1 figure and 1 table.

ASSOCIATION: Krasnodarskiy konservnyy institut (Krasnodar Institute for Canning)
Card 1/1

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

MAKASHEV, A. P.

"Carbon Dioxide as a Means of Prolonging the Storage Life of Chilled Fish Products."

Report submitted for the 10th Intl. Refrigeration Congress, Copenhagen,
19 August - 2 September 1959.

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

MAKASHEV, A. P., Doc of Tech Sci -- (diss) "Theoretical and Experimental Research on the Use of Carbon Dioxide for the Preservation of Fish and Fish Products," Krasnodar, 1959, 20 pp (Leningrad Technological Institute of the Refrigeration Industry) (KL, 2-60, 112)

MAKASHEV, A.P., kand. tekhn. nauk; MINKINA, A.I., kand. biol. nauk;
KUDAKIMOVA, A.Ya.; SOKOLOVA, Ye.V.

Effect of the intensity of proteolysis and the presence of microbes
on the occurrence of "split bellies" in some fish species. Trudy VNIRO
35:145-151 '58. (MIRA 11:11)

1. Azovskoye otdeleniye Vsesoyuznogo nauchno-issledovatel'skogo
instituta morskogo rybnogo khozyaystva i okeanografii.
(Fishery products--Preservation) (Food spoilage)

MAKASHY, A.P.; KUZNETSOV, P.G., red.; SLUZHITEL', Ye.I., tekhn.red.

[Methods of prolonging the storage of refrigerated fish]
Sposoby udlineniia srokov khraneniia okhlazhdannoi ryby.
Moskva, Vses.in-t nauchn.i tekhn.informatsii, 1958. 36 p.
(MIRA 13:4)

(Fish--Storage)

MAKASHEV, A. P.

USSR/Chemical Technology - Chemical Products and Their Application. Food Industry,
I-28

Abst Journal: Referat Zhur - Khimiya, No 19, 1956, 63722

Author: Karnitskaya, N. V., Firsova, V. I., Makashev, A. P., Aldakimova, A. Ya.

Institution: ~~None~~ A-U *Sea Res Inst. Fish Ind. & Oceanography*

Title: Action of Carbon Dioxide on Botulism Microbe in Fish Processed by Hot Smoking

Original
Periodical: Vopr. pitaniya, 1956, ^{Vol. 15} No 2, 49-50

Abstract: Study of the effects of storage of fish, that has been hot-smoked, in an atmosphere of CO₂ (70-90%) on toxin formation by B. botulinus, the spores of which are found in the intestines of some fish under natural conditions. It was found that hot-smoked fish of small and medium size is preserved in CO₂ in good condition (according to organoleptic characteristics) for 15 days as compared with 2-3 days of the controls. Storage of fish in an atmosphere of CO₂ neither inhibits nor stimulates germination of spores and toxin production of B. botulinus.

Card 1/1

MAKASHEV, A. P., BEREZIN, N. T.

Fisheries

Separation of ruff from herring by a hydraulic process. Ryb. Khoz., 28 no. 3, 1952.

MONTHLY LIST OF RUSSIAN ACCESSIONS. Library of Congress, July 1952. UNCLASSIFIED.

MAKAS, I.K., inzhener.

Rewinding sections with mica fiber casing for high-power electrical machines. Vest.elektroprom. 18 no.10:23-24 0 '47. (MLBA 6:12)

1. Energoremontrest MEP.

(Electric machinery--Maintenance and repair)

MAKARYUNAS, K. V.; MAKARYUNENE, E. K.; DARACHYUNAS, A. I. 2

"Automatic Calculation of Accidental Coincidences in Schemes for Measurement of Angular Correlations of Cascade Gamma-Rays."

report submitted for All-Union Conf on Nuclear Spectroscopy, Tbilisi, 14-22 Feb 64.

IFM LitSSR (Inst Physics & Mathematics, AS LitSSR)

MAKARYUNAS, K. V. [Makariunas, K.]

Concerning the mechanism of the reaction $\text{Be}^9(\alpha, n) \text{C}^{12}$ when
 $E_\alpha = 3,22 \text{ MeV}$. Liet sk darbai B no.1:125-127 '61. (EEAI 10:9)

1. Institut fiziki i matematiki Akademii nauk Litovskoy SSR.

(Beryllium) (Particles) (Carbon)

MAKARYUNAS, K. V. [Makariunas, K.]

On angular distribution of 12 MeV energy protons, scattered non-elastically around lithium nuclei and on spin of the lithium⁷ nucleus of 4,61 MeV energy level. Liet ak darbai B no.1:121-123 (EEAI 10:9)
'61.

1. Institut fiziki i matematiki Akademii nauk Litovskoy SSR.

(Lithium) (Protons) (Nuclear spin)

MAKARYUNAS, K. V. [Makariunas, K.]

Effective sections of some reactions (α, α) , (α, t) , (α, d) and (α, p) , flowing through lithium nuclei. Liet ak darbai B no.1: 117-120 '61. (EEAI 10:9)

L. Institut fiziki i matematiki Akademii nauk Litovskoy SSR.

(Lithium) (Particles)

33668

S/058/61/000/012/023/083
A058/A101

24.6600

AUTHORS: Starodubtsev, S. V., Makaryunas, K. V.

TITLE: Elastic and inelastic scattering of 13.2 Mev alpha particles by lithium, and the $\text{Li}^6(\alpha, p) \text{Be}^9$ and $\text{Li}^7(\alpha, p) \text{Be}^{10}$ reactions

PERIODICAL: Referativnyy zhurnal, Fizika, no. 12, 1961, 116, abstract 12B613 ("Tr. Tashkentsk. konferentsii po mirn. ispol'zovaniyu atomn. energii", 1959, v. 1, Tashkent, AN UzSSR, 1961, 98-103)

TEXT: The $\text{Li}^6(\alpha, p)$ and $\text{Li}^7(\alpha, p)$ reactions and the scattering of 13.2 Mev alpha particles by lithium were studied by the photographic emulsion method. The angular distribution of alpha particles elastically scattered by Li^7 differs from the Coulomb distribution. Proton groups from $\text{Li}^6(\alpha, p)$ and $\text{Li}^7(\alpha, p)$ reactions can be separated only for large angles. The spectrum of the proton tracks from these reactions that can be observed at $\theta_{\text{lab}} = 170^\circ$ is given. The intensity ratio of proton groups from $\text{Li}^7(\alpha, p)$ and $\text{Li}^6(\alpha, p)$ equals 1.7 ± 1 . The differential cross sections of the same groups equal 0.16 ± 0.06 and 1.17 ± 0.47 mbarn/sterad respectively. The angular distribution of protons from these two reactions are analyzed on the basis of the Butler theory.

[Abstracter's note: Complete translation]

Card 1/1

33667

3/058/61/000/012/022/083

A058/A101

Investigation of alpha-particle ...

part take place without production of composite nuclei. What is more, the authors hold that the great cross-section magnitude of these reactions (tens of millibarns) can be explained by the existence of a deuteron and a trion substructure in the Li^6 and Li^7 nuclei respectively.

[Abstracter's note: Complete translation]

X

Card 2/2

33667
S/058/61/000/012/022/083
A058/A101

24.6600

AUTHOR: Makaryunas, K. V.

TITLE: Investigation of alpha-particle interaction with lithium nuclei

PERIODICAL: Referativnyy zhurnal, Fizika, no. 12, 1961, 115-116, abstract 12B612
("Tr. Tashkentsk. konferentsii po mirn. ispol'zovaniyu atomn. energii", 1959, v. 1, Tashkent, AN UzSSR, 1961, 85-89)

TEXT: In order to study the mechanism of alpha-particle interaction with Li nuclei by the thick nuclear photoemulsion method, the angular distributions of particles from the following reactions between 10, 15, 11.5 and 13.2 Mev alpha particles and Li^6 and Li^7 were investigated: $Li^7(\alpha, \alpha') Li^7$, $Q = -4.61$ Mev; $Li^6(\alpha, d) Be^8$, $Q = -1.59$ Mev; $Li^7(\alpha, t) Be^8$, $Q = -2.56$ Mev; $Li^6(\alpha, p) Be^9$, $Q = -2.13$ Mev and $Li^7(\alpha, p) Be^{10}$, $Q = -2.56$ Mev. For the last two reactions a general angular distribution was plotted inasmuch as protons from these reactions have very close energies and could not be separated. All the angular distributions evince anisotropic structure, and their shape depends rather weakly on the energy of the bombarding particles. The form of the angular distributions indicates that $Li^6(\alpha, d)$ and $Li^7(\alpha, d)$ reactions for the most

Card 1/2

Investigation of the Reactions (α, α') , (α, p) , and (α, t) on Lithium Nuclei S/056/60/038/02/09/061
B006/B011

ASSOCIATION: Leningradskiy fiziko-tekhnicheskii institut Akademii nauk SSSR (Leningrad Institute of Physics and Technology of the Academy of Sciences, USSR)

SUBMITTED: August 7, 1959

W

Investigation of the Reactions (α, α') , (α, p) , and (α, t) on Lithium Nuclei S/056/82012/60/038/02/09/061
B006/B011

at $E_\alpha = 10.15, 11.5, \text{ and } 13.2 \text{ Mev}$, and Fig. 4 the angular distribution of protons in the center-of-mass system at $E_\alpha = 11.5 \text{ Mev}$. The angular distribution in the center-of-mass system is strongly anisotropic and asymmetric with respect to $\theta = 90^\circ$. The angular distribution of tritons originating from the reaction $\text{Li}^7(\alpha, t)\text{Be}^8$ ($Q = -2.56 \text{ Mev}$) is shown for $E_\alpha = 10.15 \text{ Mev}$ in Fig. 5, and also, for comparison, the distribution curve calculated according to Butler. A curve calculated according to the stripping theory is shown as well. It is very similar to the one of the knock-out theory. The authors finally thank the cyclotron team headed by A. B. Girshin, and also the collaborators of the laboratoriya yadernykh reaktsiy LFTI (Laboratory of Nuclear Reactions of the LFTI) for their assistance in the experiments. There are 5 figures and 14 references: 2 Soviet, 9 American, 1 British, 1 Japanese, and 1 Polish.

UH

Card 3/4

Investigation of the Reactions (α, α') , (α, p) , and (α, t) on Lithium Nuclei 82012
S/056/60/038/02/09/061
B006/B011

particle energy spectra as well as the angular distributions of the particle groups were determined. The deviations of the absolute values of the differential cross sections from the mean values did not exceed 30-40% in the various experiments. Results concerning the angular distributions of the various reactions are outlined in the paper under review. Angular distribution of reaction $\text{Li}^7(\alpha, \alpha')\text{Li}^{7*}$ ($Q = -4.61$ Mev): Fig. 2 shows the angular distribution of α -particles undergoing inelastic scattering on Li^7 , at $E_\alpha = 13.2$ Mev. The cross section calculated from an integration of the angular distribution from 15 to 90° (in the center-of-mass system) was found to be 147 ± 60 mb. A comparison (Fig. 2) with Butler's theory (Ref. 3) shows that the parity of the 4.61-Mev level of the Li^7 nucleus is negative, and that it has a spin of $1/2$, $3/2$, $5/2$, or $7/2$ (ground state of $3/2^-$). Angular distributions of the reactions $\text{Li}^6(\alpha, p)\text{Be}^9$ ($Q = -2.13$ Mev) and $\text{Li}^7(\alpha, p)\text{Be}^{10}$ ($Q = -2.56$ Mev): Fig. 3 shows the angular distributions of protons originating from these reactions in the laboratory system

Card 2/4

MAKARYUNAS, K. V.

82012
S/056/60/058/02/09/061
B006/B011

24.6600

AUTHORS: Makaryunas, K. V., Starodubtsev, S. V.TITLE: Investigation of the Reactions (α, α'), (α, p), and (α, t)
on Lithium Nuclei 19PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 38, No. 2, pp. 372 - 378

TEXT: The investigations dealt with in the present paper were conducted on the cyclotron of the Leningradskiy fiziko-tekhnicheskii institut (Leningrad Institute of Physics and Technology). α -particles with 10.15, 11.5, and 13.2 Mev were used for the experiments. A scattering chamber of 50 cm diameter was connected to the cyclotron, and the target was placed in its center; this was surrounded by photographic plates contained in special boxes. The plates were of the type Я-2 (Ya-2) with an emulsion thickness of 100 μ . The target consisted of metallic lithium in natural isotopic composition (0.75 - 1.1 mg/cm²) and was situated in dry carbon dioxide. The plates were evaluated by means of a microscope of the type МБИ-3 (MBI-3); the track lengths were measured, and the

Card 1/4

MAKARYUNAS, K. V., Cand Phys-Math Sci -- (diss) "Investigation of the reactions (a, α), (a, p), (a, d) and (a, t) with lithium nuclei." Leningrad, 1960. 10 pp; (Leningrad Physics-Technology Inst of the Academy of Sciences USSR); 250 copies; price not given; bibliography on page 10 (10 entries); (KL, 17-60, 139)

On the Mechanism of Direct Interaction in the Reaction $\text{Li}^6(\alpha, d)\text{Be}^8$ 66452
SOV/20-129-3-20/70

bombarding particles. All this, and the rather large reaction cross section are indicative of the important part played by the process developing without the formation of a compound nucleus. Probably, the α -particles knock out deuterons from the Li^6 -particles and a substructure in form of a deuteron probably exists in the Li^6 -nucleus for a certain time. There is a certain agreement between Butler's theory and experimental results. The authors thank the co-workers of the Physico-technical Institute of the AS USSR, who collaborated in the present investigation. There are 1 figure and 9 references, 2 of which are Soviet.

SUBMITTED: June 26, 1959

Card 3/3

66452

On the Mechanism of Direct Interaction in the Reaction $\text{Li}^6(\alpha, d)\text{Be}^8$ SOV/20-129-3-20/70

in the reaction $\text{Li}^6(\alpha, d)\text{Be}^8$, $Q = -1.59$ Mev. The authors investigated the angular distribution of this group. At certain small angles this group cannot be separated from the very intense group of recoil protons, so that the total angular distribution at these angles was not found. A diagram shows the angular distributions measured in the center of mass system at various energies of the bombarding α -particles (10; 15; 11.5 and 13.2 Mev). The absolute value of the

differential cross section of the reaction $\text{Li}^6(\alpha, d)\text{Be}^8$ for the angle 58° amounts to 6.7 millibarn/steradian in the center of mass system. The integral cross section approximately determined with respect to the angular distribution is, with $E_\alpha = 10.15$ Mev, not less than 50 millibarn. With an energy

increase from 10.15 to 13.2 Mev, the cross section becomes smaller. Angular distributions are sharply anisotropic, they vary continuously with a variation of energy, and are asymmetric with respect to the angle of 90° . The minimum is shifted towards smaller angles with an increasing energy of the

Card 2/3

~~21 (7)~~ 24.6600

66452

AUTHORS: Starodubtsev, S. V., Academician of the SOV/20-129-3-20/70
Uzbekskaya SSR, Makaryunas, K. V.

TITLE: On the Mechanism of Direct Interaction in the Reaction $\text{Li}^6(\alpha, d)\text{Be}^8$

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 129, Nr 3, pp 547-549 (USSR)

ABSTRACT: First, a brief report is given about the present stage of the problem on the basis of some earlier papers. The authors carried out experiments with α -particles of the energy of 8.34 and 13.2 Mev, which were accelerated in the cyclotron of the Fiziko-tekhnicheskiy institut Akademii nauk SSSR (Physico-technical Institute of the Academy of Sciences, USSR). Lithium targets of natural isotopic composition were then bombarded herewith. The particles emitted from the target were recorded on photographic plates of the type Ya-2 (emulsion thickness $100\ \mu$), which were located in a scattering chamber constructed by S. V. Starodubtsev, Ye. M. Lobanov and I. M. Shcheglov. The average angle between the plane of the photographic plate and the direction of motion of the secondary particles leaving the targets amounted to 10° . During investigation of the photographic plates under the microscope, an intense group of deuterons was found which had been produced

Card 1/3

The Angular Distributions of Tritons From the Reaction $\text{Li}^7(\alpha, t)\text{Be}^9$ SOV/56-36-5-61/76

α -particles indicates the important part played by the direct interaction mechanism. A comparison with Butler's theory shows good agreement for $l = 1$ between the theoretical and the experimental curve. The absolute values of the differential cross sections at 16° (center of mass system) are given as amounting to $9.2^{+3.7}_{-1.85}$ mb/steradian ($E_\alpha = 13.2$ Mev) and $9.4^{+4.0}_{-2.0}$ mb/steradian ($E_\alpha = 14.7$ Mev). There are 2 figures and 2 references.

ASSOCIATION: Leningradskiy fiziko-tekhnicheskii institut (Leningrad Physico-Technical Institute)

SUBMITTED: February 4, 1959

Card 2/2

21(7)
 AUTHORS: Starodubtsev, S. V., Makaryunas, K. V. SOV/56-36-5-61/76

TITLE: The Angular Distributions of Tritons From the Reaction
 $\text{Li}^7(\alpha, t)\text{Be}^8$ (Uglovyye raspredeleniya tritonov iz reaktsii
 $\text{Li}^7(\alpha, t)\text{Be}^8$)

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959,
 Vol 36, Nr 5, pp 1594-1595 (USSR)

ABSTRACT: In order to obtain information concerning the reaction
 mechanism the authors of the present "Letter to the Editor"
 investigated the angular distributions of tritons in the
 aforementioned reaction at $Q = -2.56$ Mev by the method of
 the thick photoemulsion. The α -particles were accelerated
 on the cyclotron to 8.34, 10.15, 11.5, 13.2 and 14.7 Mev. At
 all these energies similar angular distributions were
 obtained. The curves obtained representing the dependence
 of the differential cross section (in relative units) on
 the angle in the center of mass system is shown for
 $E_\alpha = 14.7$ Mev by the upper and for $E_\alpha = 13.2$ Mev by the
 lower figure. The form of the angular distributions and
 their weak dependence on the energy of the bombarding

Card 1/2

MAKARYINA, L.A. (CONTINUED)

The results agree with the calculated data obtained by the Monte Carlo method, taking into account the effect of the medium on Bremsstrahlung (Landau-Pomeranchuk and Ter-Mikaelyan effects).

For 10 cascades with $E = 1.8 \times 10^{11}$ ev, the probability of $p(\chi^2)$ from the criterion χ^2 , is $2.5 \pm 5\%$ when compared with the curves which do not consider the effect of the medium, and 80-95% when compared with the calculations that take into consideration the effect of the medium on the Bremsstrahlung.

report presented at the International Cosmic Ray Conference, Moscow 6-11 July 1959

MAKARYINA, L.A.

"ELECTRON-PHOTON CASCADES WITH ENERGIES FROM 10^{11} to 10^{13} ev IN NUCLEAR EMULSIONS"

L. S. Makaryina, A. A. Verfolomeyev, R.I. Gerasimova, I.I. Gurevich, A.S. Romantseva, S.A. Chuyeva,

Fifteen electron-photon cascades with energies from 10^{11} to 10^{13} ev, recorded in six emulsion stacks with a total volume of 10 l, have been investigated.

The energies of the primary photons evoking the cascades were determined by the energy spectrum of the cascade electrons at a depth of $2.5 - 3t_0$ (t_0 - rad. unit).

The grain density and the gap density were measured for the first pairs. In all the pairs with energies 3×10^{11} ev, a decrease in grain density at the apex caused by the screening effect was discovered.

The following experimental relation of the ionization losses of pair (1) was obtained:

where I_{pe} is the specific ionization electron loss at the ionization plateau, x is the distance from the apex of the pair in cm, and E , is the energy of the photon which produced the pair.

The expression obtained for $I/2I_{pe}$ may be used to determine the E energy from experimental values for I . An estimation of the E error is given, taking into consideration the screening effect.

The number of electron-positron pairs produced at depths of $1.0t_0$ and $1.5t_0$ was measured.

Makaryina, L.A.

"DIRECT PRODUCTION OF ELECTRON-POSITRON PAIRS BY HIGH ENERGY ELECTRONS"

L.A. Makaryina, Ap.P. Mishakova, A.S. Romantseva, G.S. Stolyarova, V.A. Turanyan, S.A. Chuyeva, A.A. Varfolomeyev, R.I. Gerasimova,

The Cross-section of direct production of electron-positron pairs by high energy electrons was measured experimentally. For this purpose, a study was made of isolated electron-photon cascades and the photon component of high energy nuclear interactions in emulsion stacks exposed to radiation in the stratosphere. In order to exclude spurious cases of direct pair production, which constitute the main difficulty in experimental measurement of the cross-section of such pairs, the calculation was carried out by the Monte Carlo method.

The calculation was made for three values of primary electron energy: 10, 100 and 1,000 Bev, taking into consideration two possible variants of the Bremsstrahlung spectrum: Bethe-Heitler and Migdal variants (Landau-Pomeranchuk and Ter-Mikaelyan effects). A method for determining the energy of ultra-relativistic electrons from the lateral distribution of the apexes of electron-positron pairs is suggested.

During the experimental measurement of very high electron energies, certain possible sources of underestimation were eliminated.

The cross section of direct pair production by high energy electrons was found to be in agreement with Ehabha's calculation within the limits of experimental error,

report presented at the International Cosmic Ray Conference, Moscow 6-11 July 1959

ILLEGIBLE

KAZANTSEV, Anatoliy Mikhaylovich, kand. tekhn. nauk, dots; Prinimali
uchastnye: LIVSHITS, I.M., inzh.; MAKAR'TEVSKIY, D.P., inzh.;
GUSEV, M.N., kand. tekhn. nauk, dotsent, retsenzent;
SHEVALDYSHEV, L.G., inzh., retsenzent; BARIT, G.Yu., red.;
VOLCHOK, K.M., tekhn. red.

[Technical norms in shipbuilding and ship repairs] Tekhnicheskoe
normirovanie v sudostroenii i sudoremonte. Leningrad, Izd-vo
"Rechnoi transport," 1962. 383 p. (MIRA 15:5)
(Shipbuilding--Production standards)
(Ships--Maintenance and repair--Production standards)

LIVSHITS, I.; MAKAR'YEVSKIY, D. P.

Setting consolidated work norms in mechanical treatment of sur-
faces, Sots.trud no.3:81-87 Mr '58. (MIRA 13:3)
(Turning--Production Standards)

MAKAR'YEVSKIY, D.P., inzhener.

Improved technology for finishing operations in ship repairs.
Resh.transp. 14 no.12:29-30 D '55. (MLRA 9:3)
(Ships--Maintenance and repair)

CHEKMAREV, A.P., akademik; KOVALENKO, Yu.Ye., kand. tekhn. nauk;
RYABOKON', N.K., inzh.; STARCHENETSkiY, M.I., inzh.;
KLYUKIN, A.R., inzh.; FOSHCHIN, A.G., inzh.; MAZAYEV, S.A.,
inzh.; BOCHKAREV, V.A., inzh.; SEZENIN, G.F.; TRAKHAN, M.I.

Investigating the process of rolling wheels at the Nizhniy
Tagil metallurgical combine. Stal' 25 no.6:543-546. Se '65.
(MIRA 12:6)

1. VNITI i Nizhne-Tagil'skiy metallurgicheskiy kombinat.

MAKARYEVA, S. P.

USSR/Chemistry - Physical chemistry

Card 1/1 : Pub. 22 - 36/46

Authors : Shigorin, D. N; Mikhaylov, N. V.; and Makaryeva, S. P.

Title : The physical structure of synthetic polyamides investigated by the infrared absorption spectra method


Periodical : Dok. AN SSSR 97/4, 711-714, Aug 1, 1954

Abstract : The application of the infrared absorption spectra method for the physico-chemical study of synthetic polyamides, is discussed. A comparison of absorption spectra of various synthetic polyamides showed that the structure of the latter is determined by a combination of three (alpha, beta, gamma) H-bonds. The existence of the three basic H-bonds in synthetic polyamides, which in fact determine their physical structure and chemical properties, was positively established. These three H-bonds are also responsible for the crystalline lattice of the synthetic polyamides. Eight references: 4-USSR and 4-USA (1936-1954). Table; drawing.

Institution : All-Union Scientific-Research Institute of Synthetic Fibers

Presented by : Academician V. A. Kargin, May 10, 1954

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6



APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

MAKAR'YEVA, S. P.

USSR/Chemistry -- Synthetic Fibers

Dec 52

"Investigation of the Physical Structure of Synthetic Polyamides by the Method of Oscillation Spectra," N. V. Mikhaylov, D. N. Shigorin and S. P. Makar'yeva, All Union Sci Res Inst of Synthetic Fibers

"Dan SSSR" Vol 87, No 6, pp 1009-1012

The nature of bonds holding the fibers together in a polycaprolactam fiber were studied using infra-red absorption spectroscopy. Comparison of the results obtained from spectral analysis with data on the mech properties of polycaprolactam confirm the assumption concerning the chain-cyclic molecular structure based on the presence of intermolecular and intramolecular hydrogen bonds. Presented by Acad A. N. Terenin 24 Oct 52

PA 240T8

MAKAR'YEVA, S. P.

FDD PA 169T20

USSR/Chemistry - Synthetic fibers, Analysis

Sep 50

"Potentiometric Determination of Sulfates in the Precipitating Bath of the Viscose Silk Manufacturing Process," S. G. Zelikman, S. P. Makar'yeva, A. B. Pakshver, All-Union Sci Res Inst of Synthetic Fiber

"Zavod Lab" Vol XVI, No 9, pp 1053-1057.

Develops method for potentiometric titration of precipitating baths with Ba chloride in presence of H peroxide. Demonstrates possibility of potentiometric titration of precipitating baths with Pb nitrate with ferroferricyanide electrode as indicator. Recommends ~~24~~ method as more efficient.

PA 169T20.

USSR/Chemistry - Chlorine
Chemistry - Analysis

Nov 1947

"Amperometric Method of Determining Chlorine," S. P. Makar'yeva, Z. G. Bezubik, M. A. Prokurinin, *Physicochemical Institute imeni L. Ya. Karpov*, 43 pp

"Zavodskaya Laboratoriya" Vol XIII, No 11

Discusses the experiments which were conducted to determine a method for measurements by amperometric calculations. Describes the experiment and lists the results. Authors were able to develop a gas analyzer for chlorine which made use of a silver electrode, and was sensitive to concentrations of 0.05 to 1.50% chlorine. They also established a proportion between

LC

36R12

Nov 1947

USSR/Chemistry - Chlorine (Contd)

the concentration of chlorine in the gaseous mixture to the strength of the depolarization current. When temperatures were raised to 40° and chlorine concentrations were 0.7% or more it was found that the depolarization current fell.

LC

36R12

MAKAR'YEVA, S. P.

MAKAR'YEVA, S. F.

Metal-plating Lab. Colloid-Electrochemical Inst.,
Acad. of Sci., USSR (-1941-)

"The Influence of Surface-active Substances and
Inorganic Salts on the Orientation of Crystals of
Copper Plating." Zhur. Fiz. Khim., Vol. 17, No. 3,
1943.

PR-52059019

PROCESSING AND PROPERTIES INDEX																									
131 AND 130 ORDERS													131 AND 130 ORDERS												
<p><i>m</i></p> <p>*The Effect of Current Density on Crystal Structure and Cathodic Polarization in the Production of Electrolytic Nickel (Coatings). S. E. Makarevich. <i>Izv. Akad. Nauk S.S.S.R. (Bull. Acad. Sci. U.R.S.S.), 1941, [Chim.], (4/5), 573-580.</i> - [In Russian.] Cf. <i>ibid.</i>, 1938, 1211; <i>Met. Abs.</i>, 1939, 6, 471. M. studied the effect of current density in the deposition of nickel on cathodic polarization and on the crystal structure, grain-size, and reflectivity of the deposits. Deposits about 20 μ thick were obtained from a solution containing 1.0M-NiSO₄, 0.2M-KCl, and 0.6M-H₂BO₃, $p_H \sim 5.0$. The c.d. was varied between 0.1 and 5.0 amp./dm.², the cathodes were of polished copper, and most of the experiments were carried out at 20° C. The results show that: (1) An increase in c.d. is accompanied by a change in orientation of the crystallites such that the order of indices of the texture axes decreases; thus, at c.d. = 0.1 amp./dm.² the axes are [113][112], at 0.5, [100][112], and at 0.9-5.0, [100][110]. At 60° C. similar changes take place, but at higher c.d.s. Replacing the stationary cathodes by rotating ones has no effect on orientation. (2) On increasing the c.d., the grain-size of the deposits increases from 1.78 μ at 0.1 amp./dm.² to a maximum of 1.97 μ at 1.5, and then diminishes to 1.41 μ at 5.0 amp./dm.². (3) The cathodic polarization rises with increase of c.d., values at 20° C. being 359 m.v. at 0.1 amp./dm.², 547 at 0.9, 635 at 1.8, 789 at 3.0, and 890 at 5.0. (4) The coeff. of reflectivity for light increases from 65% for deposits obtained at 0.1 amp./dm.² to a maximum of 90% at 2.0 amp./dm.²; there is then a decrease and the coeff. is only 80% at 5.0 amp./dm.². - N. B. V.</p>																									
<p>450-31A METALLURGICAL LITERATURE CLASSIFICATION</p>																									

CA

4

EFFECT OF TRIVALENT CHROMIUM ON THE KINETICS OF CHROMIUM PLATING. N. D. Biryukov, S. P. Makar'eva and N. D. Podobed. *Korrozija i Borba s Net* 6, No. 2, 10-14 (1940); cf. C. A. 33, 9151¹.—Two types of trivalent Cr compds. were studied (1) by introducing $\text{Cr}(\text{OH})_3$ into the electrolyte, and (2) by cathodic reduction of CrO_3 . Results were quite different. Addn. of $\text{Cr}(\text{OH})_3$ reduces the evolution of H_2 and decreases the current yield of Cr. The cathodic reduction of CrO_3 to Cr^{III} increases the evolution of H_2 , and decreases the rate of reaction at the anode. As the concn. of Cr^{III} increases, the current yield of Cr increases up to a certain max., and then decreases. For Cr^{III} from $\text{Cr}(\text{OH})_3$, the rate of anodic oxidation is 4 to 5 times that of the Cr^{III} obtained by cathodic reduction of Cr^{VI} .

C. S. Shapiro

ASB-SLA DETALLING LITERATURE CLASSIFICATION

SEARCHED INDEXED
SERIALIZED FILED
MAY 1941
NATIONAL BUREAU OF STANDARDS
WASHINGTON, D.C.

Effect of temperature on the deposition of chromium from chromic acid solutions. N. D. Biryukov and S. P. Makar'eva. *J. Applied Chem.* (U. S. S. R.) 12, 818-25 (in French, 825) (1939); cf. *C. A.* 30, 2501⁹. The Liebreich theory (cf. *C. A.* 15, 1656) of Cr plating is further developed. In the cathodic reduction of H_2CrO_4 , the velocities of the purely electrochem. reduction $\text{CrO}_4 \rightarrow \text{CrO}$ by at. H and of the oxidation of CrO by the electrolyte according to $3\text{CrO} + \text{CrO}_4 = 2\text{Cr}_2\text{O}_3$ are important. The relation between these velocities can be expressed by the equation $(C_1 \times 100)/(C_2) = K$, where the C_1 and C_2 are the percentage yields (on current) of Cr and Cr_2O_3 respectively. The value of K according to the 2nd and 1st equations. The value of K increases with an increase of temp. of Cr plating. The curve K vs. temp. of electrolyte has two points of inflection (A and B) which are characteristic for a given current. The point A (at 30°) corresponds to a sharp change in appearance and dimension of the Cr crystals. The Cr deposits below A were dull grayish and the grain size about 2.2μ ; above A the deposit is lustrous and the grain size 6.9μ . At the point B (approx. 45°) the deposits had a mirror-like appearance. The Haring and Barrows diagram (cf. *C. A.* 21, 3315) is explained by an application of A and B crit. points between which the luster deposits were obtained. A. A. Polgorny

ASH 31.4 METALLURGICAL LITERATURE CLASSIFICATION

137 AND 138 INDEX		PROCESSING AND PROPERTY INDEX		139 AND 140 INDEX	
B.C.		<p>Influence of composition of solution, current density, and temperature on properties of electrolytic nickel. S. MAKAREVA (Bull. Acad. Sci. U.R.S.S., 1933, 84r. Chim., 1911-1934).--The hardness of electrolytic Ni depends chiefly on c.d. and temp. at which deposited, whilst its coeff. of light reflexion is also influenced by the composition of the electrolyte. Ni deposited from Cl⁻ solution at 20° is considerably harder than that from SO₄²⁻ solution, but is darker in colour, porous, and not lustrous. More regular, lustrous, and harder deposits of more perfect texture are obtained from SO₄²⁻ solution containing K⁺, NH₄⁺, and Mg²⁺, i.e., cations which raise the cathode polarization of Ni. SO₄²⁻ solutions containing KCl or KF give the least porous deposits. Increase of c.d. from 0.1 to 5 amp./sq. dm. at 20° decreases the porosity and grain size and increases the hardness and lustre. At 5 amp./sq. dm. the influence of the composition of the solution disappears at 30°, but at 60° is similar to that described. Rise in temp. decreases the lustre, hardness, porosity, and perfection of texture. At 60° increase in c.d. increases grain size and decreases hardness. F. H.</p>		D-I-6	
<p>ASSOCIATE METALLURGICAL LITERATURE CLASSIFICATION</p> <p>137 AND 138 INDEX</p> <p>139 AND 140 INDEX</p>					

PROCESSING AND PROPERTY INDEX

4

ch

Theory and practice of chromium plating. III. N. D. Biryukov and S. B. Makareva. *Vestnik Metalloprod.* 14, 174-80 (1964); *Chemie & Technologie* 33, 1385; cf. C. A. 29, 1717.—Cathodic reduction of Cr^{VI} yields first Cr^{III} and then Cr^{II} , either in pure chromic acid or in presence of H_2SO_4 . The difference between the 2 cases consists in that, in pure H_2CrO_4 soln. the reduction products formed, Cr^{III} and Cr^{II} (Cr mono- and dichromate) with chromic and perchromic acids, while in presence of H_2SO_4 there is obtained sol. Cr sulfate chromate in which the proportion of H_2SO_4 is partially proportional to the H_2SO_4 concn. of the bath. At high temp. and with high c. d. H_2CrO_4 is reduced even in absence of SO_4 ions. The cause of the impossibility of obtaining a normal deposit of Cr on Pt and other metals resides in the fact that the process of reduction of Cr^{VI} is subject to the oxidizing action of chromic acid; in order to avoid it, a protective layer of a sol. chromic salt must be formed at the cathode. The sulfate chromate which forms in presence of H_2SO_4 is converted, with high H_2SO_4 concn., into $\text{Cr}_2(\text{SO}_4)_3$; the latter is reduced to Cr^{II} , which partially passes into soln. and is oxidized by the chromic acid; thus the intermediate protective layer is formed; part is also reduced to metallic Cr. In this case the electrolyte next the cathode is composed of 3 layers: the internal layer consists mainly of CrO and the intermediate layer of Cr^{III} . The latter diffuses toward the cathode where it is reduced; the CrO thus formed in turn diffuses into the intermediate layer, as does the CrO_3 from the electrolyte. In its reduction the CrO_3 oxidizes CrO to Cr_2O_3 . During the deposition of the Cr, the CrO_3 is reduced chemically, not cathodically; it therefore does not come into contact with the electrode. H_2SO_4 plays a twofold part: in its presence reduction to Cr^{II} is facilitated through the formation of sol. $\text{Cr}_2(\text{SO}_4)_3$ and the increased soly. of the CrO facilitates its reverse oxidation into Cr_2O_3 .

A. Papineau-Couture

17

ASM-SLA METALLURGICAL LITERATURE CLASSIFICATION

STANDARD REF. NO. 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 25

COMMON ELEMENTS																										PROCESS AND PROPERTIES INDEX																									
1ST AND 2ND LETTERS																										3RD AND 4TH LETTERS																									
<p><i>Ca</i></p>																										<p>Diffusion of electrolytic hydrogen through metallic palladium. S. Makarewicz. <i>J. Phys. Chem.</i> (U. S. S. R.) 8, 1380-80(1934).—At low c. ds., 0.004 amp./sq. cm. at 20-5°, all the H evolved may diffuse through a 0.085-mm. Pd cathode when the concn. of oxidizer on the other side is more than 0.05 N $K_2Cr_2O_7$, KIO_4 or $K_3Fe(CN)_6$. When the current was increased to 0.1 amp./sq. cm. only 10% diffused through. The velocity of diffusion is 6 greater for a stronger oxidizing agent but always has an upper limit regardless of increase of c. d. F. H. R.</p>																									
<p>AS H-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>																										<p>ALPHABETIC INDEX</p>																									
<p>1ST AND 2ND LETTERS</p>																										<p>3RD AND 4TH LETTERS</p>																									

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

Photoreaction of carbon tetrachloride with dioxane. Zhur.ob.khim.
31. no.12:4057-4058 D '61. (MIRA 15:2)
(Carbon tetrachloride)
(Dioxane)

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

MAKAR'YEVA, A.G.; SKUL'SKAYA, R.V.

~~Mechanization of minor operations for cold gluing of shaped parts.~~
Ier.prom.5 no.8:21 Ag '56. (MLRA 9:10)

1.Tallinskaya fanerno-mebel'naya fabrika.
(Tallinn--Furniture industry) (Gluing)

MIKONI, V.V., inzh.; LELYAVINA, L.F., tekhnik; MAKAR'YEVA, A.A., tekhnik;
VERINA, G.P., tekhn.red.

[Catalog of standard-gauge switch boxes and crossings] Al'bom
tipovykh strelochnykh perevodov i peresechenii normal'noi kolei.
No.6 [Blind crossings made of rails of the R50 and R43 types]
Glukhie peresecheniia iz rel'sov tipov R50 i R43. Moskva, Gos.
transp. zhel-dor. izd-vo. 1958. 163 p. (MIRA 12:2)

1. Moscow. Gosudarstvennyy institut tekhniko-ekonomicheskikh
izyskaniy i proyektirovaniya zheleznodorozhnogo transporta.
2. Otdel tekhnicheskikh usloviy i norm Giprottransi Ministerstva
putey soobshcheniya (for Mikoni, Lelyavina, Makar'yeva).
(Railroads--Switches)

MAKAR'YEVA, A. A.

MIKONI, V.V., inzhener; SAPRYGINA, G.M., inzhener; LELYAVINA, L.F., teknik;
MAKAR'YEVA, A.A., teknik; VERINA, T.P. tekhnicheskii redaktor.

[Album of switch boxes for normal gauge shuntings and crossings]
Al'bum tipovykh strelochnykh perevodov i peresechenii normal'noi
kolei. Moskva, Gos.transp.zhel-dor.izd-vo. Pt.2. [Ordinary switch
boxes using type R50 rails with 1/11 and 1/9 frogs] Obyknovennyye
strelochnye perevody iz rel'sov tipa R50 s krestovinyami marok
1/11 i 1/9. 1957. 172 p. (MLRA 10:6)

1. Moscow. Gosudarstvennyi institut tekhniko-ekonomicheskikh
izyskaniy i proyektirovaniya zheleznodorozhnogo transporta.
(Railroads--Switches)

MAKAR'YEV, Ye.I., inzh.

Experience in operating the Gorkiy locks. Rech.transp. 18 no.2:41-43
F '59. (MIRA 12:4)

1. Nachal'nik Gor'kovskogo shlyuza.
(Gorkiy hydroelectric power station)
(Locks (Hydraulic engineering))

MAKAR'YEV, V.V.; GALAKTIONOVA, Ye., red.; MESHCHERYAKOVA, V., tekhn.red.

[Plastics; collected texts in English] Plastmassy; sbornik
tekstov na angliiskom iazyke. Podbor tekstov, kommentarii i
slovar' V.V.Makar'eva. Moskva, Izd-vo lit-ry na inostr.iazykakh,
1960. 164 p. (MIRA 14:1)


(Plastics)

DRONIN, A.P.; ZAMANOV, V.V.; KRICHKO, A.A.; LOZOVY, A.V.; MAKAR'YEV, S.V.;
MEZHLUMOVA, A.I.; PAL'CHIKOV, G.F.; STEPURO, S.I.

Combined arrangement for the use of thermal-cracking kerosine.
Khim. i tekhn. topl. i masel 9 no.6:18-24 Je'64 (MIRA 17:7)

1. Giprogrozneft', Institut goryuchikh iskopayemykh AN SSSR i
Grozneftekhimzavody.

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6



[illegible]

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

KAZANSKIY, B.A.; DOROGUCHINSKIY, A.Z.; ROZENGART, M.I.; LYUTER, A.V.;
MITROFANOV, M.G.; BRESHCHENKO, Ye.M.; KALITA, L.A.; GOL'DSHTEYN,
Yu.A.; AFANAS'YEV, A.I.; MAKAR'YEV, S.V.; ZAMANOV, V.V.

Dehydrocyclization of normal hexane. Trudy GrozNII no. 15:
254-264 '63. (MIRA 16:5)

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

MITROFANOV, M.G.; MIRSKIY, Ya.V.; DOROGCHINSKIY, A.Z.; DRONIN, A.P.
MAKAR'YEV, S.V.; LUGOVOY, B.I.

Selecting the best arrangement for separating gasoline fractions
in molecular sieves. Trudy GrozNII no. 15:84-92 '63.
(MIRA 17:5)

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

MITROFANOV, M.G.; BONDARENKO, N.I.; MAKAR'YEV, S.V.

Technological process of dewaxing diesel fuels with crystal
carbide. Trudy GrozNII no. 15:137-142 '63. (MIRA 17:5)

A process of thermal dealkylation... S/065/62/000/004/001/004
E075/E156

pressure in benzene column 0.1-0.3 kg/cm²; temperature in benzene column, top 87 °C, bottom 130 °C; pressure in the recycle stock separation column 0.1-0.3 kg/cm²; temperature in the recycle stock separation column, top 260°, bottom 304 °C; molar ratio hydrogen/feedstock 4:1; space velocity of feed 4.0 h⁻¹; consumption of hydrogen 2.1% wt of feedstock; yield of benzene 78.7% wt of toluene. It was calculated that high grade benzene produced by the process from petroleum derived toluene is considerably cheaper than that obtained currently in the coking industry. It was established that thermal demethylation of methyl naphthalenes (700 °C, 50 atm) gives naphthalene with a yield of ca. 50% wt of feedstock after one cycle. The most suitable raw materials for the process are aromatic products obtained during reforming, pyrolysis and catalytic cracking processes. It is expected that the dealkylation process will constitute an important source of benzene and naphthalene for the Soviet petro-chemical industry. There are 1 figure and 1 table.

Card 2/2

MAKAR'YEV, S.V.

5

S/065/62/000/004/001/004
E075/E136

AUTHORS: Gonikberg, M.G., Dorogochinskiy, A.Z.,
Nitrofanov, M.G., Gavrilova, A.Ya., Dronin, A.P.,
Kupriyanov, V.A., Makar'yev, S.V., Zamanov, V.V.,
and Vovk, L.M.

TITLE: A process of thermal dealkylation of aromatic
hydrocarbons

PERIODICAL: Khimiya i tekhnologiya topliv i masel,
no.4, 1962, 11-15

TEXT: As a result of investigations carried out in the
years 1953-1960 in IOKh AN SSSR and GrozNII, a technological
scheme was developed for an industrial process of thermal
dealkylation of monocyclic aromatics such as toluene and methyl-
naphthalenes. A pilot plant for the process producing
30 000 tons of benzene per annum consists of a small number of
simple units. It contains a tubular furnace of only
3 mil. cal/hour capacity. The main production indices for the
plant are as follows: reactor pressure 50 atm; maximum
temperature 790 °C; separator temperature 35 °C;
Card 1/2

Combined Plant for the Processing of Petroleum

SOV/65-59-4-3/14

and consumption figures listed on p 16. Large-scale automation will reduce the number of operators from 54 to 14. This scheme makes it possible to manufacture high quality motor fuels, to decrease the consumption of water, steam, fuel and power. From comparative data on the efficiency of the various plants (in table 2) it is obvious that the operation of combined plants will make it possible to lower capital expenditure by 35-40% and consumption of materials by 30-45%. There is 1 figure and 2 tables.

Card 3/3

SOV/65-59-4-3/14

Combined Plant for the Processing of Petroleum

destructive distillation of goudron, the fractionation of gases with simultaneous stabilisation of petrol and the combined preparation of petrols and diesel fuels. The construction of the Giprogrozneft plant is described in detail (viz Fig). The vacuum distillate is led into the separator, after heating, and separated into the gaseous and liquid phase. The gas is used for further processing together with the catalytic cracking gas; the unstable gasoline is subjected to catalytic cracking; the fraction 195 to 380°C is mixed with the diesel fuel fraction obtained during catalytic cracking and this mixture is further processed; the fraction >380°C mixed with an analogous fraction from one of the columns, is used as a component for boiler fuels; the pitch from the evaporator DP is used either as a component of boiler fuels or utilised as raw material in coking plants. Two combined plants with an annual capacity of 3 million tons and 6 million tons of petroleum are to be based on these designs; the first plant to be used for the stabilisation of petroleum. The basic parameters of the plants are tabulated (table 1)

Card 2/3

AUTHOR: Makar'yev, S.V. SOV/65-59-4-3/14

TITLE: Combined Plant for the Processing of Petroleum
(Kombinirovannyye ustanovki dlya pererabotki nefi)

PERIODICAL: Khimiya i tekhnologiya topliv i masel, 1959, Nr 4,
pp 12-18 (USSR)

ABSTRACT: According to plans outlined at the 21st Party Congress, the extraction of petroleum is to be increased to 230-240 million tons per year by 1965; this means an annual increase of 16-18 million tons and will require the erection of new refineries. The concept of combining various processes in one plant originated in 1931. The following processes are used in the USA, Canada and other countries of the West: atmospheric-vacuum distillation, catalytic cracking and catalytic polymerisation, alkylation, catalytic reforming, catalytic and chemical purification processes etc. In the USSR the Giprogrozneft', in collaboration with GrozNII, have worked on the design of combined plants for atmospheric-vacuum distillation of petroleum, catalytic cracking of vacuum distillates, the

Card 1/3

SOV/81-59-16-58493

The Combination of Technological Processes and Installations in Oil Refining

penditures by 40% and the operation costs by 33%. The authors regard an increase in the capacity of the combined installation up to 6 million tons per year as possible.

S. Rozenoyer.

Card 2/2

SOV/81-59-16-58493

Translation from: Referativnyy zhurnal. Khimiya, 1959, Nr 16, p 409 (USSR)

AUTHORS: Strazh, A.G., Makar'yev, S.V.

TITLE: The Combination of Technological Processes and Installations in Oil Refining

PERIODICAL: Vestn. Sovnarkhoza Checheno-ingushetii, 1958, Nr 5, pp 8-11

ABSTRACT: Giprogrozneft' and GrozNII have designed a combined installation for the refining of 3 million tons of petroleum per year in which the processes of atmospheric petroleum distillation, the destructive distillation of mazut, the catalytic cracking of distillates and the head fractionation of gases are combined. For utilizing the heat obtained in the burning of coke in the regenerator of catalytic cracking, heating coils are installed through which petroleum is pumped. The heat of waste flows is also broadly used. According to the data of Giprogrozneft' the combination of the four processes in a single installation, as compared to four enlarged installations for the separate processes, reduces the capital ex-

Card 1/2

IVANOV, Anatoliy Vasil'yevich; FEDOROV, Lev Aleksandrovich;
MAKAR'YEV, P.N., red.

[Control of the engineering state and use of construction equipment in the Main Administration of the Construction Industry in Leningrad] Kontrol' za tekhnicheskim sostoianiem i ispol'zovaniem stroitel'nykh mashin v Glavleningradstroe. Leningrad, 1964, 17 p.
(MIRA 18:1)

MAKAR'YEV, P.N., inzh.

~~Utilize building machinery better. Mekh. stroi. 20 no.11:10-12 N '63.~~
(MIRA 17:1)

MAKAR'YEV, P.

New form for calculations of the work of construction equipment..
Na stroi.Ros. 3 no.9:6-7 S '62. (MIRA 15:12)

1. Glavnyy mekhanik Glavnogo stroitel'nogo upravleniya pri
ispolnitel'nom komitete Leningradskogo gorodskogo soveta
deputatov trudyashchikhsya.
(Construction equipment)

MAKAR'YEV, P.N., inzh.

Over-all mechanization and automation at the construction site.
Mekh.stroi. 18 no.9:9-12 S '61. (MIRA 14:10)

1. Glavnoye Leningradskoye upravleniye po zhilishchnomu i
grazhdanskomu stroitel'stvu.
(Building machinery)

MAKAR'YEV, P.N., inzh.

Repair of building machines by the Main Leningrad Construction Trust.
Mekh. stroi. 18 no.4:25-26 Ap '61. (MIRA 14:6)

1. Glavleningradstroy.
(Leningrad--Building machinery--Repairing)

GORBACHEV, A.I., inzh.; MAKAR'YEV, P.N., inzh.; NEFED'YEV, P.I.,
inzh.

Modernization of the SBK-1 tower crane. Mekh. stroi. 17 no.6:
12-14 Je '60. (MIRA 13:6)
(Cranes, derricks, etc.)

MAKAR'YEV, F.N.; SIROTA, M.M.; VERESOV, V.Ya., inzh., nauchnyy red.;
ROYTENBERG, A.S., red.izd-va; ROZOV, L.K., tekhn.red.

[What's new in the mechanization of construction] Novoe v mekha-
nizatsii stroitel'stva. Leningrad, Gos.izd-vo lit-ry po stroit.,
arkhit. i stroit,materialam, 1959. 62 p. (MIRA 13:6)
(Building machinery)

MAKAR'YEV, P.G.

GERSHANOV, S.V.; MAKAR'YEV, P.G.; VOL'FOVSKAYA, V.N., redaktor;
PETRUSHKO, Ye.I., tekhnicheskiy redaktor.

[Progressive practices in tractor repairing] Peredovoi opyt
remonta traktorov. Moskva, Gos. izd-vo sel'skokhoz. lit-ry,
1954. 101 p. (MLRA 7:12)
(Tractors--Repairing)

MAKAR'YEV, P. ^{G.}₂ ENG.

MAKAR'YEV, P., ENG.

Machine-Tractor Stations

Technical standardization of MTS. shops.,
MTS 12, no. 6, 1952

Monthly List of Russian Accessions, Library of Congress, October 1952. UNCLASSIFIED.

MAKAR'YEV, P., inzh.

Overall mechanization of housing construction. Na stroi. Kos.
no.11:13-15 N '61. (MIRA 16:7)

1. Glavnoye Leningradskoye upravleniye po zhilishchnomu i
grazhdanskomu stroitel'stvu.
(Construction equipment industry)

KRASNIK, F.I.; MAKAR'YEV, G.S.; SHVEDSKAYA, A.G.

Materials on the characteristics of a skin allergic test
conducted with a Rickettsia prowazekii antigen. Trudy Len.
inst. epid. i mikrobiol. 25:14-25 '63. (MIRA 17:1)

1. Iz otdela osobo opasnykh infektsiy Leningradskogo in-
stituta epidemiologii i mikrobiologii imeni Pastera i Voenno-
meditsinskoy ordena Lenina akademii imeni S.M. Kirova.

MAKAR'YEV, F. A.

"The Clinical and Laboratory Study of the Characteristics of Neurohormonal Regulation During Various Courses of Birth." Dr Med Sci, Joint Council of a Group of Leningrad Insts, Acad Med Sci USSR, Leningrad, 1954. (RZhBiol, No 3, Feb 55)

SO: Sum. No. 631, 26 Aug 55 - Survey of Scientific and Technical Dissertations Defended at USSR Higher Educational Institutions (14)

MAKAR'YEV, F. A.

MAKAR'YEV, F. A. -- "Characteristics of Neurohumoral Control During
Various Courses of Birth From a Clinical Laboratory Viewpoint." Sub 31
Oct 52, Acad Med Sci USSR. (Dissertation for the Degree of Candidate
in Medical Sciences.)

SO: Vechernaya Moskva January-December 1952

MAKAR'YEV, F. A.

"Data Concerning the Characteristics of the Cerebro-spinal
Fluid During Pregnancy and Births." Sub 16 Mar 51, Acad Med Sci USSR.

Dissertations presented for science and engineering degrees in
Moscow during 1951.

SO: Sum. No. 480, 9 May 55

15000003
ACCESSION NO. AP5007260

Analysis by the method of harmonic linearization shows that such nonlinear control laws can be found which materially enhance the noise immunity of the system to the harmonic noise. Orig. ser. has 4 figures and 50 formulas.

ASSOCIATION none

SUBJECT: CONTROL

ENCL: 00

SUB CODE: DF, IE

NO REF SOV: 003

OTHER: 000

6-4 2/1

[illegible]

Author: Andrey V. A. (Leningrad); Makarov A. B. (Leningrad)

Enhancing the noise immunity of a certain class of automatic systems by means of nonlinear control laws

1965, No. 1, Tekhnicheskaya kibernetika, no. 1, 1965, 143-149

0001 TAGS: automatic control, automatic control design, automatic control
 system, automatic control theory, noise immunity

problem is solved theoretically. Given the maximum desirable output of the nonlinear element, find the stability conditions of the system during a noise $\delta = \delta_m \sin(\Omega t + \varphi)$; the stability is to be ensured by a nonlinear control law. Two external variables are applied to the nonlinear element whose variation is given by $y(t) = A \sin(\Omega t + \varphi) + B \sin(\Omega t + \varphi) \sin(\Omega t)$, where $f_2(x)$ is a slow-varying variable and $\sin(\Omega t) = \sin(\Omega t)$ is a vibration-type noise. The

APPROVED FOR RELEASE: 06/23/11: CIA-RDP86-00513R001031500023-6

BARAN'EV, B.M.; DZERZHISKII, V.I.; PEROVSKI, B.B.,
retsensent; PETROV, F.S., red.

[Theory of remote control and homing guidance of rockets]
Teoriia sistem teleupravleniia i samonavedeniia raket. Mo-
skva, Izd-vo "Nauka," 1964. 536 p. (MIRA 17:6)